


Peru

DEMOGRAPHICS


Total population (000)	29,988	(2012)
Total under-five population (000)	2,925	(2012)
Births (000)	600	(2012)
Birth registration (%)	96	(2012)
Total under-five deaths (000)	11	(2012)
Neonatal deaths: % of all under-5 deaths	51	(2012)
Neonatal mortality rate (per 1000 live births)	9	(2012)
Infant mortality rate (per 1000 live births)	14	(2012)
Stillbirth rate (per 1000 total births)	10	(2009)
Total maternal deaths	530	(2013)
Lifetime risk of maternal death (1 in N)	440	(2013)
Total fertility rate (per woman)	2.5	(2012)
Adolescent birth rate (per 1000 girls)	67	(2010)

Under-five mortality rate
Deaths per 1000 live births



Source: IGME 2013

Maternal mortality ratio
Deaths per 100,000 live births




Source: MMEIG 2014

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition


Skilled attendant at delivery

Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)




Source: UNICEF/UNAIDS/WHO 2013

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




Source: DHS 2009

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization


Percent of children immunized:
 ■ against measles
 ■ with 3 doses Hib
 ■ with 3 doses pneumococcal conjugate vaccine
 ■ with 3 doses DTP
 ■ with rotavirus vaccine



Source: WHO/UNICEF 2013

Pneumonia treatment

Percent of children <5 years with symptoms of pneumonia:
 ■ taken to appropriate health provider
 ■ receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %)	1	(2012)	Early initiation of breastfeeding (within 1 hr of birth, %)	51	(2010)
Low birthweight incidence (moderate and severe, %)	8	(2007)	Introduction of solid, semi-solid/soft foods (%)	82	(2004-2008)
			Vitamin A two dose coverage (%)	-	

Underweight and stunting prevalence

Percent of children <5 years who are moderately or severely:



Exclusive breastfeeding

Percent of infants <6 months exclusively breastfed

